

LiquiBand[®]Surgical S

The high strength Butyl-based cyanoacrylate offering the versatility of two application tips.





Better wound closure starts with a better tool



Application Guide LiquiBand[®]Surgical S

STEP ONE

- Ensure that wound is thoroughly clean and dry prior to closure.
- While holding upright, squeeze the applicator wings until a cracking sound is heard.


STEP TWO

- Invert the tip and gently squeeze wings to prime the device.
- When the tip fills with adhesive, LiquiBand® Surgical S is ready to use.

STEP THREE

- While holding the wound edges together, apply an even coat of LiquiBand[®] Surgical S over the length of the wound.
- Continue to hold the wound edges together for 10 seconds.

Relative strengths of topical cyanoacrylates and sutures¹



STEP FOUR (Optional)

- Twist the precision tip anticlockwise and pull off the precision tip
- The deer-foot tip allows for a broad application

LiquiBand[®] Surgical S

Size	Qty Per Box	Product Code	NHS SC Code
0.8g	6	LBS-S0001	FVF 014

References 1. Data on file at Advanced Medical Solutions (Plymouth) Ltd.



- **T** +44 (0) 1752 209 955
- E info@liquiband.com
- W liquiband.com

IRM 06 0273

Advanced Medical Solutions (Plymouth) Ltd, Western Wood Way, Langage Science Park, Plympton, Plymouth, Devon, PL7 5BG, U.K. © 2018. LiquiBand is a registered trademark of Advanced Medical Solutions (Plymouth) Ltd.